

## Supplemental materials

Jablonska et al., <http://www.jcb.org/cgi/content/full/jcb.200702031/DC1>

### RT-PCR

Primers used were forward (5'-GTGACCCCTGTGGTACCGAGCACCTG-3') and reverse (5'-GGTTTGCTCTGGATGT-GGGCATGG-3') directed against the Cdk2 gene and 5'-CCCGTGATATTGCTGAAGAGCTTGGCGG-3' corresponding to the neo sequence; Olig1 sense (5'-ACTCCTTCATAAAGCCCTCG-3') and antisense primer (5'-GTGGAGATGCCACTGATGGA-3'); GFAP sense (5'-ACTTAACAAATCCCTCCTTCATCC-3') and antisense primer (5'-CCCTCTCCTGTTCACTG-3'); Tuj1 sense (5'-CCTTCGATTCCCTGGTCAGCTC-3') and antisense primer (5'-TGGGCTCCGATTCCCTCGTC-3'); EGFR sense (5'-GACCCACGAGAACTAGAAATT-3') and antisense primer (5'-AGATGGCCACACTTCACATC-3'); and actin sense (5'-CGT-GGGCCGCCCTAGGCACCA-3') and antisense primer (5'-TTGGCCTTAGGGTTCAAGGGGG-3').