

Table S1. Simulation parameters

Parameter description	Simulation	Experimental result	Reference ^a
Shape of embryo			
Long axis of egg [$\times 10^{-6}$ m]	50		
Short axes of egg [$\times 10^{-6}$ m]	30		
Radius of pronucleus [$\times 10^{-6}$ m]	5		
MT dynamics			
Growth velocity of MT (V_g) [$\times 10^{-6}$ m/s] ^b	0.328	0.118–0.8	Srayko et al., 2005
Shrinkage velocity of MT (V_s) [$\times 10^{-6}$ m/s] ^b	0.537	0.157–0.537	Kozlowski et al., 2007
Catastrophe frequency of MT (f_{cat}) [1/s] ^b	0.046	0.0115–0.046	
Rescue frequency of MT (f_{res}) [1/s] ^b	0.133	0.0113–0.133	
Number of MT fibers ^b	208	–300	Srayko et al., 2005
Pulling force			
Stall force of motor (F_{stall}) [$\times 10^{-12}$ N]	1.1	0.78–1.1	
Maximum velocity of motor (V_{max}) [$\times 10^{-6}$ m/s]	2.0	1.1–2.0	
Density of motors on MT (D) [$\times 10^3/m$] ^b	5	NA	
Ratio of number of motors on anterior and posterior cortices	2:3	2:3	Grill et al., 2003
Expected number of associating motors when an MT reaches the anterior cortex [1/s] ^b	0.8	NA	
Drag force of centrosome complexes			
Viscosity of cytosol (η) [N·s/m ²]	1	0.9–1.1	Daniels et al., 2006
Stokes' radius of the centrosome-containing complexes (r) [$\times 10^{-6}$ m] ^c	10	5–15	
Model-specific parameter			
Time step [s]	0.05		

NA, not applicable.

^aOnly references that do not appear in Kimura and Onami, 2005 are shown.^bSee Materials and methods for choice of parameters.^cStokes' radius of the centrosome-containing complexes (pronucleus and mitotic spindle) was estimated such that its value was between the radius of the pronucleus/spindle and the short radius of the egg because MTs associated with the pronucleus increase the drag force on the pronucleus (Reinsch and Gönczy, 1998).

References

- Daniels, B.R., B.C. Masi, and D. Wirtz. 2006. Probing single-cell micromechanics in vivo: the microrheology of *C. elegans* developing embryos. *Biophys. J.* 90:4712–4719.
- Grill, S.W., J. Howard, E. Schaffer, E.H. Stelzer, and A.A. Hyman. 2003. The distribution of active force generators controls mitotic spindle position. *Science*. 301:518–521.
- Kimura, A., and S. Onami. 2005. Computer simulations and image processing reveal length-dependent pulling force as the primary mechanism for *C. elegans* male pronuclear migration. *Dev. Cell*. 8:765–775.
- Kozlowski, C., M. Srayko, and F. Nedelev. 2007. Cortical microtubule contacts position the spindle in *C. elegans* embryos. *Cell*. 129:499–510.
- Reinsch, S., and P. Gönczy. 1998. Mechanisms of nuclear positioning. *J. Cell Sci.* 111:2283–2295.
- Srayko, M., A. Kaya, J. Stamford, and A.A. Hyman. 2005. *Dev. Cell*. 9:223–236.