







**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (A) BMDCs were treated with MG132 (5  $\mu$ M) or ovalbumin (500  $\mu$ g/ml) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (A) BMDCs were treated with MG132 (5  $\mu$ M) or ovalbumin (500  $\mu$ g/ml) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (A) BMDCs were treated with MG132 (5  $\mu$ M) or ovalbumin (500  $\mu$ g/ml) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (A) BMDCs were treated with MG132 (5  $\mu$ M) or ovalbumin (500  $\mu$ g/ml) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (A) BMDCs were treated with MG132 (5  $\mu$ M) or ovalbumin (500  $\mu$ g/ml) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (A) BMDCs were treated with MG132 (5  $\mu$ M) or ovalbumin (500  $\mu$ g/ml) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (A) BMDCs were treated with MG132 (5  $\mu$ M) or ovalbumin (500  $\mu$ g/ml) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (B) BMDCs were pulsed with ovalbumin (500  $\mu$ g/ml) and G9668 (3  $\mu$ M) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (B) BMDCs were pulsed with ovalbumin (500  $\mu$ g/ml) and G9668 (3  $\mu$ M) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (B) BMDCs were pulsed with ovalbumin (500  $\mu$ g/ml) and G9668 (3  $\mu$ M) for indicated time periods and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (F) WT or MyD88 KO Flt3L-derived BMDCs were pulsed with ovalbumin (500  $\mu$ g/ml) or soluble CD4-Fc protein (500  $\mu$ g/ml), or treated with LPS (10  $\mu$ g/ml) for 4 hr and analyzed by IB.





**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (F) WT or MyD88 KO Flt3L-derived BMDCs were pulsed with ovalbumin (500  $\mu$ g/ml) or soluble CD4-Fc protein (500  $\mu$ g/ml), or treated with LPS (10  $\mu$ g/ml) for 4 hr and analyzed by IB.




**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ . (F)** WT or MyD88 KO Flt3L-derived BMDCs were pulsed with ovalbumin (500  $\mu$ g/ml) or soluble CD4-Fc protein (500  $\mu$ g/ml), or treated with LPS (10  $\mu$ g/ml) for 4 hr and analyzed by IB.



**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (G) MEFs and BMDCs were pulsed with ovalbumin or treated with Tg (100 nM) for 4 hr and analyzed by IB.



**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (G) MEFs and BMDCs were pulsed with ovalbumin or treated with Tg (100 nM) for 4 hr and analyzed by IB.



**Figure S1. Antigen pulsing of BMDCs activates IRE1 $\alpha$ .** (G) MEFs and BMDCs were pulsed with ovalbumin or treated with Tg (100 nM) for 4 hr and analyzed by IB.