

Contents:

The Journal of Cell Biology

Volume 112, Number 1, January 1991

- 1 Different forms of soluble cytoplasmic mRNA binding proteins and particles in *Xenopus laevis* oocytes and embryos.
M. T. Murray, G. Krohne, and W. W. Franke
- 13 Contrasting expression patterns of three members of the *myc* family of protooncogenes in the developing and adult mouse kidney.
G. Mugrauer and P. Ekblom
- 27 Localization of components involved in protein transport and processing through the yeast Golgi apparatus.
A. Franzusoff, K. Redding, J. Crosby, R. S. Fuller, and R. Schekman
- 39 Reconstitution of constitutive secretion using semi-intact cells: regulation by GTP but not calcium.
S. G. Miller and H.-P. H. Moore
- 55 Recycling of epidermal growth factor-receptor complexes in A431 cells: identification of dual pathways.
A. Sorkin, S. Krolenko, N. Kudrjavtseva, J. Lazebnik, L. Teslenko, A. Soderquist-Mangelsdorf, and N. Nikolsky
- 65 Sequence of the clathrin heavy chain from *Saccharomyces cerevisiae* and requirement of the COOH terminus for clathrin function.
S. K. Lemmon, A. Pellicena-Palle, K. Conley, and C. L. Freund
- 81 Cell- and ligand-specific dephosphorylation of acid hydrolases. Evidence that the mannose 6-phosphatase is controlled by compartmentalization.
R. Einstein and C. A. Gabel
- 95 Structure of the mitochondrial creatine kinase octamer: high-resolution shadowing and image averaging of single molecules and formation of linear filaments under specific staining conditions.
T. Schnyder, H. Gross, H. Winkler, H. M. Eppenberger, and T. Wallimann
- 103 A Ca^{2+} transport system associated with the plasma membrane of *Dictyostelium discoideum* is activated by different chemoattractant receptors.
J. L. Milne and M. B. Coukell
- 111 Lateral diffusion and retrograde movements of individual cell surface components on single motile cells observed with nanovid microscopy.
M. de Brabander, R. Nuydens, A. Ishihara, B. Holifield, K. Jacobson, H. Geerts
- 125 The ultrastructure of patch-clamped membranes: a study using high voltage electron microscopy.
A. Rukudin, M. J. Song, and F. Sachs
- 135 Dystrophin-glycoprotein complex is highly enriched in isolated skeletal muscle sarcolemma.
K. Ohlendieck, J. M. Ervasti, J. B. Snook, and K. P. Campbell
- 149 Attachment to fibronectin or vitronectin makes human neutrophil migration sensitive to alterations in cytosolic free calcium concentration.
P. W. Marks, B. Hendey, and F. R. Maxfield
- 159 Formation of hemidesmosomes in vitro by a transformed rat bladder cell line.
K. S. Riddelle, K. J. Green, and J. C. R. Jones
- 169 Receptor functions for the integrin VLA-3: fibronectin, collagen, and laminin binding are differentially influenced by ARG-GLY-ASP peptide and by divalent cations.
M. M. Elices, L. A. Urry, and M. E. Hemler

Cover picture: Three-dimensional structure of the rhesus rotavirus by cryoelectron microscopy and icosahedral image reconstruction. The surface-shaded representations were obtained by truncating the three-dimensional maps with spherical envelopes to reveal the internal structure. See related article by Yeager et al. in the June 1990 issue, pp. 2133-2144.