

Contents:

The Journal of Cell Biology

Volume 135, Number 5, December 1996

- 1195 **Active and inactive genes localize preferentially in the periphery of chromosome territories.**
A. Kurz, S. Lampel, J.E. Nickolenko, J. Bradl, A. Benner, R.M. Zirbel, T. Cremer, and P. Lichter
- 1207 **Chromosome architecture can dictate site-specific initiation of DNA replication in *Xenopus* egg extracts.**
S.J. Lawlis, S.M. Keezer, J.-R. Wu, and D.M. Gilbert
- 1219 **Phosphorylation of linker histone is associated with transcriptional activation in a normally silent nucleus.**
M.T. Sweet, K. Jones, and C.D. Allis
- 1229 **The refolding activity of the yeast heat shock proteins Ssa1 and Ssa2 defines their role in protein translocation.**
G.L. Bush and D.I. Meyer
- 1239 **A major transmembrane protein of Golgi-derived COPI-coated vesicles involved in coatamer binding.**
K. Sohn, L. Orci, M. Ravazzola, M. Amherdt, M. Bremser, F. Lottspeich, K. Fiedler, J.B. Helms, and F.T. Wieland
- 1249 **A role for phosphoinositide 3-kinase in the completion of macropinocytosis and phagocytosis by macrophages.**
N. Araki, M.T. Johnson, and J.A. Swanson
- 1261 **The isoforms of proprotein convertase PC5 are sorted to different subcellular compartments.**
I. De Bie, M. Marcinkiewicz, D. Malide, C. Lazure, K. Nakayama, M. Bendayan, and N.G. Seidah
- 1277 **Roles for Rac1 and Cdc42 in planar polarization and hair outgrowth in the wing of *Drosophila*.**
S. Eaton, R. Wepf, and K. Simons
- 1291 **F-Actin bundles in *Drosophila* bristles are assembled from modules composed of short filaments.**
L.G. Tilney, P. Connelly, S. Smith, and G.M. Guild
- 1309 **Bundling of actin filaments by elongation factor 1 α inhibits polymerization at filament ends.**
J.W. Murray, B.T. Edmonds, G. Liu, and J. Condeelis
- 1323 ***MHP1*, an essential gene in *Saccharomyces cerevisiae* required for microtubule function.**
I. Irminger-Finger, E. Hurt, A. Roebuck, M.A. Collart, and S.J. Edelstein
- 1341 **Genetic and metabolic status of NGF-deprived sympathetic neurons saved by an inhibitor of ICE family proteases.**
M. Deshmukh, J. Vasilakos, T.L. Deckwerth, P.A. Lampe, B.D. Shivers, and E.M. Johnson, Jr.
- 1355 **Molecular composition of the node of Ranvier: Identification of ankyrin-binding cell adhesion molecules neurofascin (mucin+/third FNIII domain-) and NrCAM at nodal axon segments.**
J.Q. Davis, S. Lambert, and V. Bennett
- 1369 **Presence of double-strand breaks with single-base 3' overhangs in cells undergoing apoptosis but not necrosis.**
V.V. Didenko and P.J. Hornsby
- 1377 **A role for Jun-N-terminal kinase in anoikis; suppression by bcl-2 and crmA.**
S.M. Frisch, K. Vuori, D. Kelaita, and S. Sicks
- 1383 **Inhibition of pp125^{FAK} in cultured fibroblasts results in apoptosis.**
J.E. Hungerford, M.T. Compton, M.L. Matter, B.G. Hoffstrom, and C.A. Otey
- 1391 **Translation of CUG- but not AUG-initiated forms of human fibroblast growth factor 2 is activated in transformed and stressed cells.**
S. Vagner, C. Touriol, B. Galy, S. Audigier, M.-C. Gensac, F. Amalric, F. Bayard, H. Prats, and A.-C. Prats
- 1403 **Molecular and functional defects in kidneys of mice lacking collagen α 3(IV): Implications for alport syndrome.**
J.H. Miner and J.R. Sanes
- 1415 **Reduction of type V collagen using a dominant-negative strategy alters the regulation of fibrillogenesis and results in the loss of corneal-specific fibril morphology.**
J.K. Marchant, R.A. Hahn, T.F. Linsenmayer, and D.E. Birk

Cover picture: The cover shows a scanning electron micrograph of phagosomes and ruffles on the surface of a macrophage (blue) and partially engulfed sheep erythrocytes (pink). Incubation in wortmannin, an inhibitor of phosphoinositide 3-kinase, allowed ruffling and the extension of pseudopodia, but prevented completion of the phagocytosis. The micrograph was taken by Dr. Nobukazu Araki of Ehime University (magnification 10,000). See related article by Araki et al., 1249–1260.