Contents:

The Journal of Cell Biology

Volume 137, Number 2, April 21, 1997

- 263 Components of the human SWI/SNF complex are enriched in active chromatin and are associated with the nuclear matrix.
 - J.C. Reves, C. Muchardt, and M. Yaniv
- 275 Structural protein 4.1 in the nucleus of human cells: Dynamic rearrangements during cell division.
 - S.W. Krauss, C.A. Larabell, S. Lockett, P. Gascard, S. Penman, N. Mohandas, and J.A. Chasis
- 291 The dynamic nuclear redistribution of an hnRNP K-homologous protein during *Drosophila* embryo development and heat shock. Flexibility of transcription sites in vivo.
 - P. Buchenau, H. Saumweber, and D.J. Arndt-Jovin
- Assembly and intracellular targeting of the $\beta\gamma$ subunits of heterotrimeric G proteins.
 - A. Rehm and H.L. Ploegh
- 319 Dissociation of coatomer from membranes is required for brefeldin A-induced transfer of Golgi enzymes to the endoplasmic reticulum. J. Scheel, R. Pepperkok, M. Lowe, G. Griffiths, and T.E. Kreis
- Mannose 6-phosphate receptors regulate the formation of clathrin-coated vesicles in the TGN.R. Le Borgne and B. Hoflack
- Segregation of glucosylceramide and sphingomyelin occurs in the apical to basolateral transcytotic route in HepG2 cells.
 S.C.D. van IJzendoorn, M.M.P. Zegers, J.W. Kok, and D. Hoekstra
- 359 The apical submembrane cytoskeleton participates in the organization of the apical pole in epithelial cells.
 - P.J.I. Salas, M.L. Rodriguez, A.L. Viciana, D.E. Vega-Salas, and H.-P. Hauri
- 377 Tim23, a protein import component of the mitochondrial inner membrane, is required for normal activity of the multiple conductance channel, MCC.
 - T.A. Lohret, R.E. Jensen, and K.W. Kinnally

- 387 Regulated membrane localization of Tiam1, mediated by the NH₂-terminal pleckstrin homology domain, is required for Rac-dependent membrane ruffling and c-Jun NH₂-terminal kinase activation.
 - F. Michiels, J.C. Stam, P.L. Hordijk, R.A. van der Kammen, L. Ruuls-Van Stalle, C.A. Feltkamp, and J.G. Collard
- 399 High rates of actin filament turnover in budding yeast and roles for actin in establishment and maintenance of cell polarity revealed using the actin inhibitor latrunculin-A.
 - K.R. Ayscough, J. Stryker, N. Pokala, M. Sanders, P. Crews, and D.G. Drubin
- 417 The *Saccharomyces cerevisiae* kinesin-related motor Kar3p acts at preanaphase spindle poles to limit the number and length of cytoplasmic microtubules.
 - W. Saunders, D. Hornack, V. Lengyel, and C. Deng
- 433 A role for mitogen-activated protein kinase in the spindle assembly checkpoint in XTC cells. X.M. Wang, Y. Zhai, and J.E. Ferrell, Jr.
- 445 Synaptic-like microvesicles of neuroendocrine cells originate from a novel compartment that is continuous with the plasma membrane and devoid of transferrin receptor.
 - A. Schmidt, M.J. Hannah, and W.B. Huttner
- Differentiation and death of premyelinating oligodendrocytes in developing rodent brain.
 - B.D. Trapp, A. Nishiyama, D. Cheng, and W. Macklin

Contents continued

Cover picture: Confocal image of myelinating oligodendrocyte in developing rat brain immunostained with an antibody specific for proteolipid protein (green) and another specific for an epitope common to proteolipid protein and its spliced variant, DM20 (red). Staining patterns indicate targeting of DM20 to the plasma membrane of oligodendrocyte perikarya and processes (red) and to compact myelin (yellow). In contrast, PLP appears to be targeted only to compact myelin (yellow). See related article in this issue by Trapp et al. 459–468.

469 Processing/activation of at least four interleukin-1β converting enzyme-like proteases occurs during the execution phase of apoptosis in human monocytic tumor cells.

M. MacFarlane, K. Cain, X.-M. Sun, E.S. Alnemri, and G.M. Cohen

481 Regulation of cell motility by mitogen-activated protein kinase.

R.L. Klemke, S. Cai, A.L. Giannini, P.J. Gallagher, P. de Lanerolle, and D.A. Cheresh

493 ICAMs redistributed by chemokines to cellular uropods as a mechanism for recruitment of T lymphocytes.

M.A. del Pozo, C. Cabañas, M.C. Montoya, A. Ager, P. Sánchez-Mateos, and F. Sánchez-Madrid

509 Isolated P-selectin glycoprotein ligand-1 dynamic adhesion to P- and E-selectin.

D.J. Goetz, D.M. Greif, H. Ding, R.T. Camphausen, S. Howes, K.M. Comess, K.R. Snapp, G.S. Kansas, and F.W. Luscinskas

521 ADDITIONS AND CORRECTIONS