Contents:

The Journal of Cell Biology

Volume 140, Number 4, February 23, 1998

Mini-Review

733 Toxin entry: Retrograde transport through the secretory pathway.

J.M. Lord and L.M. Roberts

Regular Articles

737 SRPK2: A differentially expressed SR protein-specific kinase involved in mediating the interaction and localization of pre-mRNA splicing factors in mammalian cells.

H.-Y. Wang, W. Lin, J.A. Dyck, J.M. Yeakley, Z. Songyang, L.C. Cantley, and X.-D. Fu

751 gp25L/emp24/p24 protein family members of the cis-Golgi network bind both COP I and II coatomer

M. Dominguez, K. Dejgaard, J. Füllekrug, S. Dahan, A. Fazel, J.-P. Paccaud, D.Y. Thomas, J.J.M. Bergeron, and T. Nilsson

767 Oligomeric structure of type I and type II transforming growth factor β receptors: Homodimers form in the ER and persist at the plasma membrane.

L. Gilboa, R.G. Wells, H.F. Lodish, and Y.I. Henis

779 A novel dynamin-like protein associates with cytoplasmic vesicles and tubules of the endoplasmic reticulum in mammalian cells.

Y. Yoon, K.R. Pitts, S. Dahan, and M.A. McNiven

795 Caveolin-1 and -2 in the exocytic pathway of MDCK cells.

P. Scheiffele, P. Verkade, A.M. Fra, H. Virta, K. Simons, and E. Ikonen

807 A mobile PTS2 receptor for peroxisomal protein import in *Pichia pastoris*.

Y. Elgersma, M. Elgersma-Hooisma, T. Wenzel, J.M. McCaffery, M.G. Farquhar, and S. Subramani

821 Rapid diffusion of green fluorescent protein in the mitochondrial matrix.

A. Partikian, B. Ölveczky, R. Swaminathan, Y. Li, and A.S. Verkman

Role of ryanodine receptors in the assembly of calcium release units in skeletal muscle.

F. Protasi, C. Franzini-Armstrong, and P.D. Allen

A transgenic myogenic cell line lacking ryanodine receptor protein for homologous expression studies: Reconstitution of Ry₁R protein and function.

R.A. Moore, H. Nguyen, J. Galceran, I.N. Pessah, and P.D. Allen

853 Titin extensibility in situ: Entropic elasticity of permanently folded and permanently unfolded molecular segments.

K. Trombitás, M. Greaser, S. Labeit, J.-P. Jin, M. Kellermayer, M. Helmes, and H. Granzier

861 Cytoskeletal polarization of T cells is regulated by an immunoreceptor tyrosine-based activation motif-dependent mechanism.

B. Lowin-Kropf, V.S. Shapiro, and A. Weiss

873 Smy1p, a kinesin-related protein that does not require microtubules.

S.H. Lillie and S.S. Brown

885 Ezrin/radixin/moesin (ERM) proteins bind to a positively charged amino acid cluster in the juxta-membrane cytoplasmic domain of CD44, CD43, and ICAM-2.

S. Yonemura, M. Hirao, Y. Doi, N. Takahashi, T. Kondo, Sa. Tsukita, and Sh. Tsukita

897 A class VI unconventional myosin is associated with a homologue of a microtubule-binding protein, cytoplasmic linker protein–170, in neurons and at the posterior pole of *Drosophila* embryos.

V.A. Lantz and K.G. Miller

911 The p75 neurotrophin receptor mediates neuronal apoptosis and is essential for naturally occurring sympathetic neuron death.

S.X. Bamji, M. Majdan, C.D. Pozniak, D.J. Belliveau, R. Aloyz, J. Kohn, C.G. Causing, and F.D. Miller

925 Disrupted proteolipid protein trafficking results in oligodendrocyte apoptosis in an animal model of Pelizaeus-Merzbacher disease.

A. Gow, C.M. Southwood, and R.A. Lazzarini

Contents continued

Cover picture: Cytoskeletal polarization of a Jurkat T cell towards an anti-TCR-coated latex bead as revealed by confocal microscopy. Focal engagement of the TCR induces the reorientation of the microtubule-organizing center towards the site of cell-bead contact, whereas F-actin-rich pseudopods surround the bead. The image shows a series of z-sections through a Jurkat T cell-bead conjugate double stained for microtubules (green) and actin (red). The microtubule-organizing center appears as a brightly stained green dot. See related article in this issue by Lowin-Kropf et al., 861–871.

- 935 Hepatocyte nuclear factor 4 provokes expression of epithelial marker genes, acting as a morphogen in dedifferentiated hepatoma cells. G.F. Späth and M.C. Weiss
- 947 Vascular endothelial growth factor induces endothelial fenestrations in vitro.
 - S. Esser, K. Wolburg, H. Wolburg, G. Breier, T. Kurzchalia, and W. Risau

- 961 CAS/Crk coupling serves as a "molecular switch" for induction of cell migration. R.L. Klemke, J. Leng, R. Molander, P.C. Brooks, K. Vuori, and D.A. Cheresh
- 973 ADDITIONS AND CORRECTIONS