## **Contents:**

## The Journal of Cell Biology

Volume 144, Number 2, January 25, 1999

## In Brief

Making peroxisomes from (almost) nothing. Multi-step nuclear envelope reassembly. Rho as a positional signal for cytokinesis. Separating guidance from fasciculation. Disulfides in the cytoplasm.

W.A. Wells

## Regular Articles

- 201 Nuclear import of the parsley bZIP transcription factor CPRF2 is regulated by phytochrome photoreceptors.
  - S. Kircher, F. Wellmer, P. Nick, A. Rügner, E. Schäfer, and K. Harter
- 213 Nuclear import of Cdk/cyclin complexes: Identification of distinct mechanisms for import of Cdk2/cyclin E and Cdc2/cyclin B1.
  - J.D. Moore, J. Yang, R. Truant, and S. Kornbluth
- 225 Temporal differences in the appearance of NEP-B78 and an LBR-like protein during *Xenopus* nuclear envelope reassembly reflect the ordered recruitment of functionally discrete vesicle types.
  - S. Drummond, P. Ferrigno, C. Lyon, J. Murphy, M. Goldberg, T. Allen, C. Smythe, and C.J. Hutchison
- 241 Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release in chromaffin cells seen from inside the ER with targeted aequorin.

M.T. Alonso, M.J. Barrero, P. Michelena, E. Carnicero, I. Cuchillo, A.G. García, J. García-Sancho, M. Montero, and J. Alvarez

- 255 Peroxisome synthesis in the absence of preexisting peroxisomes.
  - S.T. South and S.J. Gould
- An unconventional role for cytoplasmic disulfide bonds in vaccinia virus proteins.
  - J. Krijnse Locker and G. Griffiths
- 281 Ordering the cytochrome c-initiated caspase cascade: Hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner.
  - E.A. Slee, M.T. Harte, R.M. Kluck, B.B. Wolf, C.A. Casiano, D.D. Newmeyer, H.-G. Wang, J.C. Reed, D.W. Nicholson, E.S. Alnemri, D.R. Green, and S.J. Martin
- 293 Characterization of a *Chlamydomonas* insertional mutant that disrupts flagellar central pair microtubule-associated structures.
  - D.R. Mitchell and W.S. Sale

- 305 The small GTP-binding protein Rho regulates cortical activities in cultured cells during division.
  - C.B. O'Connell, S.P. Wheatley, S. Ahmed, and Y.-l. Wang
- 315 Dissociation of FAK/p130<sup>CAS</sup>/c-Src complex during mitosis: Role of mitosis-specific serine phosphorylation of FAK.
  - Y. Yamakita, G. Totsukawa, S. Yamashiro, D. Fry, X. Zhang, S.K. Hanks, and F. Matsumura
- Neural cell adhesion molecule (N-CAM) is required for cell type segregation and normal ultrastructure in pancreatic islets.
  - F. Esni, I.-B. Täljedal, A.-K. Perl, H. Cremer, G. Christofori, and H. Semb
- 339 Neurolin Ig domain 2 participates in retinal axon guidance and Ig domains 1 and 3 in fasciculation.
  - C.A. Leppert, H. Diekmann, C. Paul, U. Laessing, M. Marx, M. Bastmeyer, and C.A.O. Stuermer
- 351 Analysis of C-cadherin regulation during tissue morphogenesis with an activating antibody.
  - Y. Zhong, W.M. Brieher, and B.M. Gumbiner
- 361 The junction-associated protein AF-6 interacts and clusters with specific Eph receptor tyrosine kinases at specialized sites of cell-cell contact in the brain.
  - M. Buchert, S. Schneider, V. Meskenaite, M.T. Adams, E. Canaani, T. Baechi, K. Moelling, and C.M. Hovens
- 373 Membrane-type 1 matrix metalloprotease (MT1-MMP) enables invasive migration of glioma cells in central nervous system white matter.
  - A.T.J. Beliën, P.A. Paganetti, and M.E. Schwab

Cover picture: Computer averages of digitized thin section EM images, modified by an edge transformation, reveal the normal structure of central pair microtubule complexes from *Chlamydomonas flagella* (left). Different projections are missing from the central pair complex in flagellar motility mutants cpc1 (middle) and pf6 (right). In the wild-type both of these projections repeat at 16-nm intervals. The role of central pair structures in regulation of flagellar dynein activity is discussed in the article by Mitchell and Sale, 293–304.