Contents:

The Journal of Cell Biology

Volume 146, Number 4, August 23, 1999

In Brief

Cooperating cytoskeletons. Defining the real prereplication complex. Giant gene manipulation. Phosphoiositide control of endocytosis.

W.A. Wells

Mini-Reviews

697 Employment of the epidermal growth factor receptor in growth factor–independent signaling pathways.

G. Carpenter

703 Extranuclear apoptosis: The role of the cytoplasm in the execution phase.

J.C. Mills, N.L. Stone, and R.N. Pittman

Regular Articles

709 Mcm2, but not RPA, is a component of the mammalian early G1-phase prereplication complex.

D.S. Dimitrova, I.T. Todorov, T. Melendy, and D.M. Gilbert

723 The ribosome regulates the GTPase of the β-subunit of the signal recognition particle receptor.

G. Bacher, M. Pool, and B. Dobberstein

731 The developmental role of *warthog*, the Notch modifier encoding *Drab6*.

K. Purcell and S. Artavanis-Tsakonas

741 Electrospray ionization tandem mass spectrometry (ESI-MS/MS) analysis of the lipid molecular species composition of yeast subcellular membranes reveals acyl chain-based sorting/remodeling of distinct molecular species en route to the plasma membrane.

R. Schneiter, B. Brügger, R. Sandhoff, G. Zellnig, A. Leber, M. Lampl, K. Athenstaedt, C. Hrastnik, S. Eder, G. Daum, F. Paltauf, F.T. Wieland, and S.D. Kohlwein

755 Phosphoinositide–AP-2 interactions required for targeting to plasma membrane clathrin-coated pits.

I. Gaidarov and J.H. Keen

765 VAMP-7 mediates vesicular transport from endosomes to lysosomes.

R.J. Advani, B. Yang, R. Prekeris, K.C. Lee, J. Klumperman, and R.H. Scheller

777 Sid2p, a spindle pole body kinase that regulates the onset of cytokinesis.

C.A. Sparks, M. Morphew, and D. McCollum

791 Regulation of APC activity by phosphorylation and regulatory factors.

S. Kotani, H. Tanaka, H. Yasuda, and K. Todokoro

801 Domains in the 1α dynein heavy chain required for inner arm assembly and flagellar motility in *Chlamydomonas*.

S.H. Myster, J.A. Knott, K.M. Wysocki, E. O'Toole, and M.E. Porter

819 In vivo, villin is required for Ca²⁺-dependent F-actin disruption in intestinal brush borders.

E. Ferrary, M. Cohen-Tannoudji, G. Pehau-Arnaudet, A. Lapillonne, R. Athman, T. Ruiz, L. Boulouha, F. El Marjou, A. Doye, J.-J. Fontaine, C. Antony, C. Babinet, D. Louvard, F. Jaisser, and S. Robine

Integrating the actin and vimentin cytoskeletons: Adhesion-dependent formation of fimbrin-vimentin complexes in macrophages.

I. Correia, D. Chu, Y.-H. Chou, R.D. Goldman, and P. Matsudaira

843 Analysis of CD44-containing lipid rafts: Recruitment of annexin II and stabilization by the actin cytoskeleton.

S. Oliferenko, K. Paiha, T. Harder, V. Gerke, C. Schwärzler, H. Schwarz, H. Beug, U. Günthert, and L.A. Huber

855 Exogenous expression of β-catenin regulates contact inhibition, anchorage-independent growth, anoikis, and radiation-induced cell cycle arrest

K. Orford, C.C. Orford, and S.W. Byers

Contents continued

Cover picture: Chinese hamster ovary cells were synchronized at the onset of S-phase and the earliest-replicating DNA sequences were briefly pulse-labeled with 5-chloro-2'-deoxyuridine (CldU). Prelabeled cells were chased into the next cell cycle and synchronized during G1 (left column) or at the beginning of S-phase (right column). The recruitment of replicating proteins to these chromosomal domains was examined by immunolabeling with antibodies specific for CldU (green) and either Mcm2 (red, top row) or RPA (red, bottom row). Mcm2 was bound to early-replicating chromatin throughout G1-phase and was displaced from replicons shortly after their initiation. By contrast, RPA associated transiently with chromosomal domains at the specific time of S-phase when they replicated. See related article in this issue by Dimitrova et al., 709–722.

- Intracellular movement of green fluorescent protein-tagged phosphatidylinositol 3-kinase in response to growth factor receptor signaling.
 H. Gillham, M.C.H.M. Golding, R. Pepperkok, and W.J. Gullick
- Insulin-like growth factor II signaling in neoplastic proliferation is blocked by transgenic expression of the metalloproteinase inhibitor TIMP-1.

D.C. Martin, J.L. Fowlkes, B. Babic, and R. Khokha

- 893 Role of transmembrane 4 superfamily (TM4SF) proteins CD9 and CD81 in muscle cell fusion and myotube maintenance.
 - I. Tachibana and M.E. Hemler