

Table S1. *The Effects of Several Inflammatory Agents on Thymocytes*

Thymus	n ^a	Total	DN	DP	CD8 SP	CD4 SP
PBS	17	790 ± 101 ^b	30.5 ± 4.2	663 ± 88.3	36.8 ± 6.4	60.4 ± 7.4
NP-CGG/IFA	3	817 ± 12.5	41.7 ± 1.2	674 ± 79.3	36.5 ± 0.8	64.1 ± 7.6
TNFα	15	594 ± 54.1	29.0 ± 3.6	511 ± 47.3	21.8 ± 2.9 ^c	34.1 ± 3.6 ^d
IL-1β	7	543 ± 88.9	24.9 ± 7.9	465 ± 74.0	23.8 ± 6.2	29.5 ± 5.7 ^d
IL-6	5	841 ± 187	31.8 ± 7.1	714 ± 161	27.6 ± 2.6	67.9 ± 13.7
IFNβ	6	586 ± 194	20.7 ± 5.0	555 ± 169	31.5 ± 3.3	57.5 ± 12.8
PTX	3	775 ± 101	24.3 ± 21.6	617 ± 143	98.6 ± 42.9	98.6 ± 42.9

^aTotal numbers of mice from multiple (two to eight) experiments.

^bAverage ± SEM ($\times 10^6$) number of cells recovered from the femurs of individual mice 6 d after treatment.

^cValue is significantly different from the corresponding population in control mice, P ≤ 0.05.

^dValues are significantly different from the corresponding population in control mice, P < 0.01.