

## Siegers et al., <http://www.jem.org/cgi/content/full/jem.20070782/DC1>

### SUPPLEMENTAL MATERIALS AND METHODS

**Antibodies.** The rabbit anti- $\zeta$  antiserum 448 (9) and the anti-hCD3 $\delta$  antibody (APA1/2) (10) have been previously described. Jovi1 (anti-hC $\beta$ 1) was provided by M. Owen (GlaxoSmithKline, Herts, UK). UCHT1 (anti-hCD3 $\epsilon$ ) was provided by P. Beverly (The Edward Jenner Institute for Vaccine Research, Compton, UK). 145-2C11 (anti-mCD3 $\epsilon$ ) was provided by J. Bluestone (University of California, San Francisco, San Francisco, CA), and HMT3.2 (anti-hCD3 $\gamma$ ) was provided by R. Kubo (Gemini Science Inc., La Jolla, CA). The following reagents were purchased: 17A2 (anti-mCD3 $\gamma$ ), GL3-PE (anti-mTCR $\gamma\delta$ ), biotin-labeled 145-2C11, goat anti-mouse FITC, and streptavidin-Cy5 (BD Biosciences); 7D6 (anti-mCD3 $\gamma$ ; Biocarta Europe GmbH); 12CA5 (anti-HA tag; Roche); 5A6.E9 (anti-hTCR $\gamma\delta$ ; Pierce Chemical Co.);  $\beta$ F1 (anti-hTCR $\beta$ ; Endogen); M208 (anti-CD3 $\epsilon$ ) and M20 (anti-CD3 $\epsilon$ ; Santa Cruz Biotechnology, Inc.); G3 (anti- $\zeta$ ; Abcam); 4G10 (antiphosphotyrosine; Upstate Biotechnology); goat anti-rabbit IgG-HRPO and rabbit anti-goat IgG-HRPO (Pierce Chemical Co.); OKT3 (anti-hCD3 $\epsilon$ ; American Type Culture Collection); anti-CD4 and anti-CD8 (Immunotech); 11F2 (anti-hTCR $\gamma\delta$ ; BD Biosciences); Immu510 (anti-hTCR $\delta$ ; Beckman Coulter) and SK7 (anti-hCD3 $\epsilon$ ; StemCell Technologies Inc.); UCHT1-Alexa Fluor 647 (anti-hCD3 $\epsilon$ ; Biozol); and goat anti-mouse IgG-PE (Southern Biotechnology Associates, Inc.).

### REFERENCES

1. Rolli, V., M. Gallwitz, T. Wossning, A. Flemming, W.W. Schamel, C. Zurn, and M. Reth. 2002. Amplification of B cell antigen receptor signaling by a Syk/ITAM positive feedback loop. *Mol. Cell.* 10:1057–1069.
2. Siegers, G.M., J. Yang, C.U. Duerr, P.J. Nielsen, M. Reth, and W.W. Schamel. 2006. Identification of disulfide bonds in the Ig- $\alpha$ /Ig- $\beta$  component of the B cell antigen receptor using the *Drosophila* S2 cell reconstitution system. *Int. Immunol.* 18:1385–1396.
3. Haks, M.C., T.A. Cordaro, J.H. van den Brakel, J.B. Haanen, E.F. de Vries, J. Borst, P. Krimpenfort, and A.M. Kruisbeek. 2001. A redundant role of the CD3 gamma-immunoreceptor tyrosine-based activation motif in mature T cell function. *J. Immunol.* 166:2576–2588.
4. Call, M.E., J. Pyrdol, M. Wiedmann, and K.W. Wucherpfennig. 2002. The organizing principle in the formation of the T cell receptor-CD3 complex. *Cell.* 111:967–979.
5. Wegener, A.M., X. Hou, J. Dietrich, and C. Geisler. 1995. Distinct domains of the CD3-gamma chain are involved in surface expression and function of the T cell antigen receptor. *J. Biol. Chem.* 270:4675–4680.
6. Arnett, K.L., S.C. Harrison, and D.C. Wiley. 2004. Crystal structure of a human CD3-epsilon/delta dimer in complex with a UCHT1 single-chain antibody fragment. *Proc. Natl. Acad. Sci. USA.* 101:16268–16273.
7. Sun, Z.J., K.S. Kim, G. Wagner, and E.L. Reinherz. 2001. Mechanisms contributing to T cell receptor signaling and assembly revealed by the solution structure of an ectodomain fragment of the CD3 epsilon gamma heterodimer. *Cell.* 105:913–923.
8. Sun, Z.Y., S.T. Kim, I.C. Kim, A. Fahmy, E.L. Reinherz, and G. Wagner. 2004. Solution structure of the CD3epsilon/delta ectodomain and comparison with CD3epsilon/gamma as a basis for modeling T cell receptor topology and signaling. *Proc. Natl. Acad. Sci. USA.* 101:16867–16872.
9. Gil, D., W.W. Schamel, M. Montoya, F. Sanchez-Madrid, and B. Alarcon. 2002. Recruitment of Nck by CD3 epsilon reveals a ligand-induced conformational change essential for T cell receptor signaling and synapse formation. *Cell.* 109:901–912.
10. Alarcon, B., S.C. Ley, F. Sanchez-Madrid, R.S. Blumberg, S.T. Ju, M. Fresno, and C. Terhorst. 1991. The CD3-gamma and CD3-delta subunits of the T cell antigen receptor can be expressed within distinct functional TCR/CD3 complexes. *EMBO J.* 10:903–912.