Volume 153 No. 2 February 1, 1981

- 221 JOEL D. TAUROG, ELIZABETH S. RAVECHE, PATRICIA A. SMATHERS, LAURIE H. GLIMCHER, DAVID P. HUSTON, CARL T. HANSEN, and ALFRED D. STEINBERG. T cell abnormalities in NZB mice occur independently of autoantibody production
- 235 JAMES E. ROBINSON, DOUGLAS SMITH, and JAMES NIEDERMAN. Plasmacytic differentiation of circulating Epstein-Barr virus-infected B lymphocytes during acute infectious mononucleosis
- 245 GEORGE S. DRUMMOND and ATTALLAH KAPPAS. Potent heme-degrading action of antimony and antimony-containing parasiticidal agents
- 257 MARGARET J. POLLEY, RALPH L. NACHMAN, and BABETTE B. WEKSLER. Human complement in the arachidonic acid transformation pathway in platelets
- 269 ROBERT L. COFFMAN and IRVING L. WEISSMAN. A monoclonal antibody that recognizes B cells and B cell precursors in mice
- 280 E. J. Jenkinson, W. Van Ewijk, and J. J. T. Owen. Major histocompatibility complex antigen expression on the epithelium of the developing thymus in normal and nude mice
- 293 M. Dy, B. Lebel, P. Kamoun, and J. Hamburger. Histamine production during the antiallograft response. Demonstration of a new lymphokine enhancing histamine synthesis
- JEFFREY A. LEDBETTER, ROBERT L. EVANS, MARC LIPINSKI, CHARLOTTE CUNNINGHAM-RUNDLES, ROBERT A. GOOD, and LEONARD A. HERZENBERG. Evolutionary conservation of surface molecules that distinguish T lymphocyte helper/inducer and cytotoxic/suppressor subpopulations in mouse and man
- 324 Shozo Izui, Robert A. Eisenberg, and Frank J. Dixon. Subclass-restricted IgG polyclonal antibody production in mice injected with lipid A-rich lipopolysaccharides
- 339 KENNETH A. AULT and MURRAY TOWLE. Human B lymphocyte subsets. I. IgG-bearing B cell response to pokeweed mitogen
- 352 J. LATHAM CLAFLIN, SUSAN HUDAK, and ANNE MADDALENA. Anti-phosphocholine hybridoma antibodies. I. Direct evidence for three distinct families of antibodies in the murine response
- Wen-Yuan Chen, Jose Muñoz, H. Hugh Fudenberg, Edward Tung, and Gabriel Vi-Rella. Polyclonal activation of human peripheral blood B lymphocytes by fomaldehydefixed Salmonella paratyphi B. I. Immunoglobulin production without DNA synthesis
- 375 Masao Kimoto and C. Garrison Fathman. Antigen-reactive T cell clones. II. Unique homozygous and (high responder × low responder)F₁ hybrid antigen-presenting determinants detected using poly(Tyr, Glu)-poly D, L-Ala--poly Lys-reactive T cell clones
- 386 MARCUS A. HORWITZ and SAMUEL C. SILVERSTEIN. Interaction of the Legionnaires' disease bacterium (*Legionella pneumophila*) with human phagocytes. I. L. pneumophila resists killing by polymorphonuclear leukocytes, antibody, and complement
- 398 MARCUS A. HORWITZ AND SAMUEL C. SILVERSTEIN. Interaction of the Legionnaires' disease bacterium (*Legionella pneumophila*) with human phagocytes. II. Antibody promotes binding of *L. pneumophila* to monocytes but does not inhibit intracellular multiplication
- 407 DAVID H. KATZ, LEE R. KATZ, and CHERYL A. BOGOWITZ. Cell-interaction molecules on immunocompetent lymphocytes. Development of anti-parent cell-interaction-molecule-receptor reactions in F₁ hybrid mice and evidence for a unique F₁ hybrid subset of interacting cells
- 423 ALAIN J. DESSEIN, WENDY L. PARKER, STEPHANIE L. JAMES, and JOHN R. DAVID. IgE antibody and resistance to infection. I. Selective suppression of the IgE antibody response in rats diminishes the resistance and the eosinophil response to *Trichinella spiralis* infection
- JONATHAN S. BROMBERG, BARUJ BENACERRAF, and MARK I. GREENE. Mechanisms of regulation of cell-mediated immunity. VII. Suppressor T cells induced by suboptimal doses of antigen plus an I-J-specific allogeneic effect
- 450 MICHAEL H. DIETZ, MAN-SUN SY, BARUJ BENACERRAF, ALFRED NISONOFF, MARK I. GREENE, and RONALD N. GERMAIN. Antigen- and receptor-driven regulatory mechanisms. VII. H-2-restricted anti-idiotypic suppressor factor from efferent suppressor T cells

BRIEF DEFINITIVE REPORTS

- 464 MARY MICHAELIDES, MAURO SANDRIN, GILLIAN MORGAN, IAN F. C. McKENZIE, ROBERT ASHMAN, and ROGER W. MELVOLD. Ir gene function in an I-A subregion mutant B6.C-H-2^{bm12}
- 470 ABBY L. MAIZEL, SHASHIKANT R. MEHTA, RICHARD J. FORD, and LAWRENCE B. LACHMAN. Effect of interleukin 1 on human thymocytes and purified human T cells
- 476 Perrie B. Hausman, Daniel P. Stittes, and John D. Stobo. Antigen-reactive T cells can be activated by autologous macrophages in the absence of added antigen
- 482 EDWARD J. GOETZL and WALTER C. PICKETT. Novel structural determinants of the human neutrophil chemotactic activity of leukotriene B
- 488 HSIU-SAN LIN, SAIMON GORDON, DIEN-MING CHEN, and MICHAEL KURTZ. Conversion of monocytes to cells capable of anchorage-independent growth in vitro
- 494 ARNOLD E. POSTLETHWAITE, JORMA KESKI-OJA, GARY BALIAN, and ANDREW H. KANG. Induction of fibroblast chemotaxis by fibronectin. Localization of the chemotactic region to a 140,000-molecular weight non-gelatin-binding fragment

PREPARATION OF MANUSCRIPT Articles should conform to the style of a current issue of this journal or to the recommendations of the Style Manual for Biological Journals (4th edition 1978, American Institute of Biological Sciences, 1401 Wilson Blvd., Arlington, Va. 22209). Note that each reference should contain the title of the pertinent citation. Please supply a brief title for a running head, not exceeding 57 characters and spaces. Only original typescript is acceptable for the printer; in addition, submit a fully legible carbon, mimeographed, or Xerox copy. Double space entire manuscript, including references, legends, footnotes, and tables.

PREPARATION OF ILLUSTRATIONS Label all illustrations (photographs, micrographs, charts, or drawings) sequentially with arabic numerals; designate them as figures. Number tables with roman numerals. In the margin of the manuscript, indicate approximately where each figure or table should appear.

PAGE SIZE The text block on each Journal page is $5 \times 7\%$ inches. Plan illustrations, including their legends, to fit within these dimensions after reduction. Avoid figures that exceed 5 inches in width unless they can fill the entire page, including margins. The maximum area for such full-page figures, including their legends, is $5\frac{1}{2} \times 8$.

LINE DRAWINGS Provide original ink drawing (preferably $8\frac{1}{2} \times 11$ inches maximum). These give the sharpest reproductions. The lettering should be sans serif, and of consistent size, and so planned that the size of the lettering is in scale with the size and complexity of the figures. Lettering and labeling should be large enough to permit reduction to a minimum size. Capital letters and numbers should be $1\frac{1}{2}$ to $1\frac{3}{4}$ mm high after reduction.

MICROGRAPHS Micrographs should be planned so that they can be reproduced same size (not reduced). Figure numbers and lettering should be $2\frac{1}{2}$ to 3 mm high in sans serif style. If authors are unable to provide figures labeled satisfactorily, the Press will label them; in this case, the author must provide an overlay or diagram to show exactly where such labeling should appear.

In general, limit the field of a micrograph to the regions and structures discussed in the report; eliminate irrelevant structures or expanses. Cropping unnecessary areas from micrographs saves space and expense. If a micrograph must be specifically oriented, mark "top" on the back of the print.

Micrographs that are grouped should be of uniform size and shape, with the edges of the prints touching. The engraver will tool thin white lines between figures. Plan the over-all size of the grouping so the legend will fit below the composite; if the grouping occupies a full page, the legend will appear at the foot of the facing page. The Journals Office prefers that multiple photographs be submitted unmounted with a diagram showing the desired layout.