

# CONTENTS THE JOURNAL OF EXPERIMENTAL MEDICINE

Volume 158 No. 4 October 1, 1983

- 
- 1021 KYOKO IIDA, LEE NADLER, and VICTOR NUSSENZWEIG. Identification of the membrane receptor for the complement fragment C3d by means of a monoclonal antibody
  - 1034 CRAIG M. SORENSEN, CARL W. PIERCE, and DAVID R. WEBB. Purification and characterization of an L-glutamic acid<sup>60</sup>-L-alanine<sup>30</sup>-L-tyrosine<sup>10</sup> (GAT)-specific suppressor factor from genetic responder mice
  - 1048 GIOVANNA TOSATO, SANDRA E. PIKE, and R. MICHAEL BLAESE. Reversal of infectious mononucleosis-associated suppressor T cell activity by D-mannose
  - 1061 TERRY A. POTTER, MICHAEL A. PALLADINO, DARCY B. WILSON, and T. V. RAJAN. Epitopes on H-2D<sup>d</sup> somatic cell mutants recognized by cytotoxic T cells
  - 1077 PHILIPPA MARRACK, ROBERT ENDRES, RICHARD SHIMONKEVITZ, ALBERT ZLOTNIK, DENO DIALYNAS, FRANK FITCH, and JOHN KAPPLER. The major histocompatibility complex-restricted antigen receptor on T cells. II. Role of the L3T4 product
  - 1092 BICE PERUSSIA, ELAHE TALIEH DAYTON, ROBERT LAZARUS, VIRGINIA FANNING, and GIORGIO TRINCHIERI. Immune interferon induces the receptor for monomeric IgG1 on human monocytic and myeloid cells
  - 1114 SOMAN N. ABRAHAM, DAVID L. HASTY, W. ANDREW SIMPSON, and EDWIN H. BEACHEY. Antiadhesive properties of a quaternary structure-specific hybridoma antibody against type 1 fimbriae of *Escherichia coli*
  - 1129 LEONARD J. RUBINSTEIN, BURT GOLDBERG, JACQUES HIERNAUX, KATHRYN E. STEIN, and CONSTANTIN A. BONA. Idiotype-antiidiotype regulation. V. The requirement for immunization with antigen or monoclonal antiidiotypic antibodies for the activation of  $\beta 2 \rightarrow 6$  and  $\beta 2 \rightarrow 1$  polyfructosan-reactive clones in BALB/c mice treated at birth with minute amounts of anti-A48 idiotype antibodies
  - 1145 GILLA KAPLAN, WESLEY C. VAN VOORHIS, EUZENIR NUNES SARNO, NADIA NOGUEIRA, and ZANVIL A. COHN. The cutaneous infiltrates of leprosy: a transmission electron microscopy study
  - 1160 BRUCE N. CRONSTEIN, SARA B. KRAMER, GERALD WEISSMANN, and ROCHELLE HIRSCHHORN. Adenosine: a physiological modulator of superoxide anion generation by human neutrophils
  - 1178 YOSHIHIRO ASANO and RICHARD J. HODES. T cell regulation of B cell activation: cloned Lyt-1<sup>+</sup>2<sup>-</sup> T suppressor cells inhibit the major histocompatibility complex-restricted interaction of T helper cells with B cells and/or accessory cells
  - 1191 MASAHIRO KUROSAKA and MORRIS ZIFF. Immunoelectron microscopic study of the distribution of T cell subsets in rheumatoid synovium
  - 1211 D. E. MADDOX, J. H. BUTTERFIELD, S. J. ACKERMAN, C. B. COULAM, and G. J. GLEICH. Elevated serum levels in human pregnancy of a molecule immunochemically similar to eosinophil granule major basic protein
  - 1227 JEFFREY L. PLATT, BARBARA W. GRANT, ALLISON A. EDDY, and ALFRED F. MICHAEL. Immune cell populations in cutaneous delayed-type hypersensitivity
  - 1243 ANJANA RAO, SUSAN J. FAAS, LINDA J. MILLER, PHILLIP S. RIBACK, and HARVEY CANTOR. Lysis of inducer T cell clones by activated macrophages and macrophage-like cell lines
  - 1259 LAWRENCE Y. C. AGODOA, V. JOYCE GAUTHIER, and MART MANNIK. Precipitating antigen-antibody systems are required for the formation of subepithelial electron-dense immune deposits in rat glomeruli
  - 1272 ZENA WERB and JENNIE R. CHIN. Apoprotein E is synthesized and secreted by resident and thioglycollate-elicited macrophages but not by pyran copolymer- or bacillus Calmette-Guerin-activated macrophages
  - 1294 PATRICIA BALDACCI, FRANÇOISE POZO, SYLVIE GISSELBRECHT, and PHILIPPE KOURILSKY. Altered transcription of genes coding for class I histocompatibility antigens in murine tumor cells
  - 1307 PERRIE B. HAUSMAN, CHARLES E. MOODY, JUDITH B. INNES, JAMES J. GIBBONS, and MARC E. WEKSLER. Studies on the syngeneic mixed lymphocyte reaction. III. Development of a monoclonal antibody with specificity for autoreactive T cells

CONTENTS CONTINUED ON REVERSE OF THIS COVER

- 1319 MARCUS A. HORWITZ. Formation of a novel phagosome by the Legionnaires' disease bacterium (*Legionella pneumophila*) in human monocytes

#### BRIEF DEFINITIVE REPORTS

- 1332 RICHARD J. ROBB and WARNER C. GREENE. Direct demonstration of the identity of T cell growth factor-binding protein and the Tac antigen
- 1338 SAMUEL D. WRIGHT, LYDIA S. CRAIGMYLE, and SAMUEL C. SILVERSTEIN. Fibronectin and serum amyloid P component stimulate C3b- and C3bi-mediated phagocytosis in cultured human monocytes
- 1344 OLIVERA J. FINN, CINDY J. STACKHOUSE, and RICHARD S. METZGAR. Human Ia beta chains and the invariant chain share a common antigenic determinant
- 1350 NANCY E. ADELMAN, DAVID L. WATLING, and HUGH O. MCDEVITT. Treatment of (NZB  $\times$  NZW) $F_1$  disease with anti-I-A monoclonal antibodies
- 1356 E. A. GRIMM, R. J. ROBB, J. A. ROTH, L. M. NECKERS, L. B. LACHMAN, D. J. WILSON, and S. A. ROSENBERG. Lymphokine-activated killer cell phenomenon. III. Evidence that IL-2 is sufficient for direct activation of peripheral blood lymphocytes into lymphokine-activated killer cells
- 1362 SUBRAMANIAM SRIRAM and LAWRENCE STEINMAN. Anti-I-A antibody suppresses active encephalomyelitis: treatment model for diseases linked to IR genes
- 1368 ORESTE ACUTO, STEFAN C. MEUER, JAMES C. HODGDON, STUART F. SCHLOSSMAN, and ELLIS L. REINHERZ. Peptide variability exists within  $\alpha$  and  $\beta$  subunits of the T cell receptor for antigen
- 1374 ROBERT W. KARR, CAROL C. KANNAPELL, RICHARD J. ZANGARA, SANNA M. GOYERT, JACK SILVER, and BENJAMIN D. SCHWARTZ. An HLA-DR5 homozygous cell line expresses two DS (I-A-like) molecules
- 1381 CORRECTION

#### COPYEDITING AND PRODUCTION *Heather Libbey and Richard Cunniff*

**PREPARATION OF MANUSCRIPT** Articles should conform to the style of a current issue of this journal or to the recommendations of the *Council of Biology Editors Style Manual* (4th edition, 1978, American Institute of Biological Sciences, 1401 Wilson Blvd., Arlington, Va. 22209). Note that each reference should contain the title of the pertinent citation. Please supply a brief title for a running head, not exceeding 57 characters and spaces, with no abbreviations. Only original typescript is acceptable for the printer; in addition, submit a fully legible carbon, mimeographed, or Xerox copy. Double space entire manuscript, including references, legends, footnotes, and tables.

**PREPARATION OF ILLUSTRATIONS** Label all illustrations (photographs, micrographs, charts, or drawings) sequentially with arabic numerals; designate them as figures. Number tables with roman numerals. In the margin of the manuscript, indicate approximately where each figure or table should appear.

**PAGE SIZE** The text block on each Journal page is  $5 \times 7\frac{1}{4}$  inches. Plan illustrations, including their legends, to fit within these dimensions after reduction. Avoid figures that exceed 5 inches in width unless they can fill the entire page, including margins. The maximum area for such full-page figures, including their legends, is  $5\frac{1}{2} \times 8$ .

**LINE DRAWINGS** Provide original ink drawing (preferably  $8\frac{1}{2} \times 11$  inches maximum). These give the sharpest reproductions. The lettering should be sans serif, and of consistent size, and so planned that the size of the lettering is in scale with the size and complexity of the figures. Lettering and labeling should be large enough to permit reduction to a minimum size. Capital letters and numbers should be  $1\frac{1}{2}$  to  $1\frac{3}{4}$  mm high after reduction.

**MICROGRAPHS** Micrographs should be planned so that they can be reproduced same size (not reduced). Figure numbers and lettering should be  $2\frac{1}{2}$  to 3 mm high in sans serif style. If authors are unable to provide figures labeled satisfactorily, the Press will label them; in this case, the author must provide an overlay or diagram to show exactly where such labeling should appear.

In general, limit the field of a micrograph to the regions and structures discussed in the report; eliminate irrelevant structures or expanses. Cropping unnecessary areas from micrographs saves space and expense. If a micrograph must be specifically oriented, mark "top" on the back of the print.

Micrographs that are grouped should be of uniform size and shape, with the edges of the prints touching. The engraver will tool thin white lines between figures. Plan the over-all size of the grouping so the legend will fit below the composite; if the grouping occupies a full page, the legend will appear at the foot of the facing page. The Journals Office prefers that multiple photographs be submitted unmounted with a diagram showing the desired layout.