

CONTENTS THE JOURNAL OF EXPERIMENTAL MEDICINE

Volume 159 No. 2 February 1, 1984

-
- i HENRY G. KUNKEL, 1916-1983
 - 341 T. J. BRACIALE, V. L. BRACIALE, T. J. HENKEL, J. SAMBROOK, and M.-J. GETHING. Cytotoxic T lymphocyte recognition of the influenza hemagglutinin gene product expressed by DNA-mediated gene transfer
 - 355 JACQUES VAN SNICK, JEAN-PAUL COUTELIER, EMIEL VAN ROOST, and JEAN-LOUIS GUÉNET. Rheumatoid factors in 129XB recombinant inbred strains. Igh-1-linked control of allotypic and isotypic specificities
 - 365 URSULA KEES and PETER H. KRAMMER. Most influenza A virus-specific memory cytotoxic T lymphocytes react with antigenic epitopes associated with internal virus determinants
 - 378 ROBERTO S. ACCOLLA. Analysis of the structural heterogeneity and polymorphism of human Ia antigens. Four distinct subsets of molecules are coexpressed in the Ia pool of both DR 1,1 homozygous and DR 3,W6 heterozygous B cell lines
 - 394 GERALD T. NEPOM, BARBARA S. NEPOM, PAOLO ANTONELLI, ERIC MICKELSON, JACK SILVER, SANNA M. GOYERT, and JOHN A. HANSEN. The HLA-DR4 family of haplotypes consists of a series of distinct DR and DS molecules
 - 405 KOHKICHI YAMAMOTO and RICHARD B. JOHNSTON, JR. Dissociation of phagocytosis from stimulation of the oxidative metabolic burst in macrophages
 - 417 GÜNTHER HEINRICH, ANDRÉ TRAUNECKER, and SUSUMU TONEGAWA. Somatic mutation creates diversity in the major group of mouse immunoglobulin κ light chains
 - 436 PAMELA J. FINK, MICHAEL J. BEVAN, and IRVING L. WEISSMAN. Thymic cytotoxic T lymphocytes are primed in vivo to minor histocompatibility antigens
 - 452 M. S. BLAKE and E. C. GOTSCHLICH. Purification and partial characterization of the opacity-associated proteins of *Neisseria gonorrhoeae*
 - 463 GÉRALD J. PRUD'HOMME, ABRAHAM FUKS, ELEANOR COLLE, THOMAS A. SEEMAYER, and RONALD D. GUTTMAN. Immune dysfunction in diabetes-prone BB rats. Interleukin 2 production and other mitogen-induced responses are suppressed by activated macrophages
 - 479 ANJANA RAO, SUSAN J. FAAS, and HARVEY CANTOR. Activation specificity of arsonate-reactive T cell clones. Structural requirements for hapten recognition and comparison with monoclonal antibodies
 - 495 AMITABHA MAZUMDER and STEVEN A. ROSENBERG. Successful immunotherapy of natural killer-resistant established pulmonary melanoma metastases by the intravenous adoptive transfer of syngeneic lymphocytes activated in vitro by interleukin 2
 - 508 STEVEN T. PALS, HELGA GLEICHMANN, and ERNST GLEICHMANN. Allosuppressor and allo-helper T cells in acute and chronic graft-vs.-host disease. V. F₁ mice with secondary chronic GVHD contain F₁-reactive allohelper but no allosuppressor T cells
 - 524 W. SLUITER, INE ELZENGA-CLAASEN, ANNELIES VAN DER VOORT VAN DER KLEY-VAN ANDEL, and R. VAN FURTH. Differences in the responses of inbred mouse strains to the factor increasing monocytopoiesis
 - 537 LENNART OLSSON, RINETTE B. ANDREASEN, ÅKE OST, BIRGER CHRISTENSEN, and PETER BIBERFELD. Antibody-producing human-human hybridomas. II. Derivation and characterization of an antibody specific for human leukemia cells
 - 551 THOMAS HÜNIG. Monoclonal anti-Lyt-2.2 antibody blocks lectin-dependent cellular cytotoxicity of H-2-negative target cells
 - 559 ARMAND BENSUSSAN, STEFAN C. MEUER, STUART F. SCHLOSSMAN, and ELLIS L. REINHERZ. Delineation of an immunoregulatory amplifier population recognizing autologous Ia molecules. Analysis with human T cell clones
 - 577 E. L. KHOURY, G. F. BOTTAZZO, and I. M. ROITT. The thyroid "microsomal" antibody revisited. Its paradoxical binding in vivo to the apical surface of the follicular epithelium
 - 592 ANN AGER and JOHN L. GORDON. Differential effects of hydrogen peroxide on indices of endothelial cell function

CONTENTS CONTINUED ON REVERSE OF THIS COVER

- 604 I. F. ROWE, ANNE K. SOUTAR, IRIS M. TRAYNER, MARILYN L. BALTZ, F. C. DE BEER, LOUISE WALKER, D. BOWYER, J. HERBERT, A. FEINSTEIN, and M. B. PEPYS. Rabbit and rat C-reactive proteins bind apolipoprotein B-containing lipoproteins

BRIEF DEFINITIVE REPORTS

- 617 SHIN-ICHI NISHIKAWA, TATSUO KINA, JUN-ICHIRO GYOTOKU, and YOSHIMOTO KATSURA. High frequency of λ gene activation in bone marrow pre-B cells
- 623 MARY MCNAMARA and HEINZ KOHLER. Regulatory idiotopes: induction of idiotype-recognizing helper T cells by free light and heavy chains
- 629 DAVID WOFSY, CHRISTOPHER E. KERGER, and WILLIAM E. SEAMAN. Monocytosis in the BXSB model for systemic lupus erythematosus
- 635 KENNETH E. BERNSTEIN, EDMUNDO LAMOYI, NANCY MCCARTNEY-FRANCIS, ROSE G. MAGE. Sequence of a cDNA encoding Basilea kappa light chains (K2 isotype) suggests a possible relationship of protein structure to limited expression
- 641 JEFFREY S. HOFFMAN, LOWELL H. ERICSSON, NILS ERIKSEN, KENNETH A. WALSH, and EARL P. BENDITT. Murine tissue amyloid protein AA: NH₂-terminal sequence identity with only one of two serum amyloid protein (ApoSAA) gene products
- 647 LINDA ANDRUS, ANGELA GRANELLI-PIPERNO, and E. REICH. Cytotoxic T cells both produce and respond to interleukin 2

COPYEDITING AND PRODUCTION *Heather Libbey and Richard Cunniff*

PREPARATION OF MANUSCRIPT Articles should conform to the style of a current issue of this journal or to the recommendations of the *Council of Biology Editors Style Manual* (4th edition, 1978, American Institute of Biological Sciences, 1401 Wilson Blvd., Arlington, Va. 22209). Note that each reference should contain the title of the pertinent citation. Please supply a brief title for a running head, not exceeding 57 characters and spaces, with no abbreviations. Capitalize trade names and give manufacturers' full names and addresses. Only original typescript is acceptable for the printer; in addition, submit a fully legible carbon, mimeographed, or Xerox copy. Double space entire manuscript, including references, legends, footnotes, and tables.

PREPARATION OF ILLUSTRATIONS Label all illustrations (photographs, micrographs, charts, or drawings) sequentially with arabic numerals; designate them as figures. Number tables with roman numerals. In the margin of the manuscript, indicate approximately where each figure or table should appear.

PAGE SIZE The text block on each Journal page is 5 × 7¼ inches. Plan illustrations, including their legends, to fit within these dimensions after reduction.

LINE DRAWINGS Provide original ink drawing (preferably 8½ × 11 inches maximum). These give the sharpest reproductions. The lettering should be sans serif, and of consistent size, and so planned that the size of the lettering is in scale with the size and complexity of the figures. Lettering and labeling should be large enough to permit reduction to a minimum size. Capital letters and numbers should be 1½ to 1¾ mm high after reduction.

MICROGRAPHS Micrographs should be planned so that they can be reproduced same size (not reduced). Figure numbers and lettering should be 2½ to 3 mm high in sans serif style. If authors are unable to provide figures labeled satisfactorily, the Press will label them; in this case, the author must provide an overlay or diagram to show exactly where such labeling should appear.

In general, limit the field of a micrograph to the regions and structures discussed in the report; eliminate irrelevant structures or expanses. Cropping unnecessary areas from micrographs saves space and expense. If a micrograph must be specifically oriented, mark "top" on the back of the print.

Micrographs that are grouped should be of uniform size and shape, with the edges of the prints touching. The engraver will tool thin white lines between figures. Plan the over-all size of the grouping so the legend will fit below the composite; if the grouping occupies a full page, the legend will appear at the foot of the facing page. The Journals Office prefers that multiple photographs be submitted unmounted with a diagram showing the desired layout.