## Volume 159 No. 4 April 1, 1984

- 971 MASATAKA KASAI, TOHRU TAKASHI, TOSHITADA TAKAHASHI, and TOHRU TOKUNAGA. A new differentiation antigen (FT-1) shared with fetal thymocytes and leukemic cells in the mouse
- 981 YOSHINARI TAKASAKI, DIANE FISHWILD, and ENG M. TAN. Characterization of proliferating cell nuclear antigen recognized by autoantibodies in lupus sera
- 993 JOSEPH H. PHILLIPS, AN MY LE, and LEWIS L. LANIER. Natural killer cells activated in a human mixed lymphocyte response culture identified by expression of Leu-11 and class II histocompatibility antigens
- 1009 ERIC G. NEILSON, EILEEN MCCAFFERTY, S. MICHAEL PHILLIPS, MICHAEL D. CLAYMAN, and CAROLYN J. KELLY. Antiidiotypic immunity in interstitial nephritis. II. Rats developing anti-tubular basement membrane disease fail to make an antiidiotypic regulatory response: the modulatory role of an RT7.1<sup>+</sup>, OX8<sup>-</sup> suppressor T cell mechanism
- 1027 D. W. GOLDMAN and E. J. GOETZL. Heterogeneity of human polymorphonuclear leukocyte receptors for leukotriene B<sub>4</sub>. Identification of a subset of high affinity receptors that transduce the chemotactic response
- 1042 BETTY P. TSAO, DARYL S. FAIR, LINDA K. CURTISS, and THOMAS S. EDGINGTON. Monocytes can be induced by lipopolysaccharide-triggered T lymphocytes to express functional factor VII/VIIa protease activity
- 1058 C. R. K. HIND, P. M. COLLINS, D. RENN, R. B. COOK, D. CASPI, MARILYN L. BALTZ, and M. B. PEPYS. Binding specificity of serum amyloid P component for the pyruvate acetal of galactose
- 1070 GEORGE DRIZLIKH, JOSEPHA SCHMIDT-SOLE, and BORIS YANKELEVICH. Involvement of the K and I regions of the H-2 complex in resistance to hemopoietic allografts
- 1083 V. A. FISCHETTI, K. F. JONES, B. N. MANJULA, and J. R. SCOTT. Streptococcal M6 protein expressed in *Escherichia coli*. Localization, purification, and comparison with streptococcalderived M protein
- 1096 MASARU TANIGUCHI, TAKESHI TOKUHISA, TOSHIHIRO ITOH, and MASAMOTO KANNO.

  Functional roles of two polypeptide chains that compose an antigen-specific suppressor T cell factor
- 1105 Masatoshi Tateno, Nobuo Kondo, Tetsuo Itoh, Tsugihiko Chubachi, Takehiro Togashi, and Takashi Yoshiki. Rat lymphoid cell lines with human T cell leukemia virus production. I. Biological and serological characterization
- 1117 THOMAS J. PALKER, RICHARD M. SCEARCE, SARA E. MILLER, MIKOLAS POPOVIC, DANI P. BOLOGNESI, ROBERT C. GALLO, and BARTON F. HAYNES. Monoclonal antibodies against human T cell leukemia-lymphoma virus (HTLV) p24 internal core protein. Use as diagnostic probes and cellular localization of HTLV
- 1132 JOHN P. DALEY and ICHIRO NAKAMURA. Natural resistance of lethally irradiated F<sub>1</sub> hybrid mice to parental marrow grafts is a function of H-2/Hh-restricted effectors
- 1149 BARTON F. HAYNES, RICHARD M. SCEARCE, DAVID F. LOBACH, and LUCINDA L. HEN-SLEY. Phenotypic characterization and ontogeny of mesodermal-derived and endocrine epithelial components of the human thymic microenvironment
- 1169 R. R. HARDY, K. HAYAKAWA, D. R. PARKS, L. A. HERZENBERG, and L. A. HERZENBERG. Murine B cell differentiation lineages
- 1192 DAVID H. SHERR and MARTIN E. DORF. An idiotype-specific helper population that bears immunoglobulin, Ia, and Lyt-1 determinants
- 1201 ELLEN WHITNACK, JAMES B. DALE, and EDWIN H. BEACHEY. Common protective antigens of group A streptococcal proteins masked by fibrinogen
- 1213 Julia L. Greenstein, John Kappler, Philippa Marrack, and Steven J. Burakoff. The role of L3T4 in recognition of Ia by a cytotoxic, H-2D<sup>d</sup>-specific T cell hybridoma
- 1225 ROBIN E. CALLARD, CARRIE M. WINGER, and SHANE L. TIERNAN. Antigen-specific suppression of human antibody responses by allogeneic T cells. I. Frequency and antigen specificity of allogeneic suppressor T cells and their role in major histocompatibility complex-controlled genetic restriction

- 1238 KENNETH L. ROCK and BARUJ BENACERRAF. Selective modification of a private I-A allo stimulating determinant(s) upon association of antigen with an antigen-presenting cell
- 1253 DANIÉLÉ PRIMI and PIERRE-ANDRÉ CAZENAVE. The B cell repertoire revealed by major histocompatibility complex-specific helper T cells. I. Frequencies of a genetically defined V region marker among mitogen- and T helper cell-reactive B lymphocytes in normal and immunized mice
- 1270 GERARDO R. VASTA, JOHN J. MARCHALONIS, and HEINZ KOHLER. Invertebrate recognition protein cross-reacts with an immunoglobulin idiotype

## BRIEF DEFINITIVE REPORTS

- 1277 HARUMI JYONOUCHI and PAUL W. KINCADE. Precocious and enhanced functional maturation of B lineage cells in New Zealand black mice during embryonic development
- 1283 THOMAS L. ROTHSTEIN, RICHARD A. MILLER, DEBRA J. PARKER, ENID KELLY, A. PAUL VASTOLA, and ANN MARSHAK-ROTHSTEIN. Induction of idiotope suppression in the anti-azophenylarsonate response of T-depleted A/J mice
- 1289 LIANA SCHLESINGER, ROBERT A. MUSSON, and RICHARD B. JOHNSON, JR. Functional and biochemical studies of multinucleated giant cells derived from the culture of human monocytes

## COPYEDITING AND PRODUCTION Heather Libbey and Richard Cunniff

PREPARATION OF MANUSCRIPT Articles should conform to the style of a current issue of this journal or to the recommendations of the Council of Biology Editors Style Manual (4th edition, 1978, American Institute of Biological Sciences, 1401 Wilson Blvd., Arlington, Va. 22209). Note that each reference should contain the title of the pertinent citation. Please supply a brief title for a running head, not exceeding 55 characters and spaces, with no abbreviations. Capitalize trade names and give manufacturers' full names and addresses. Only original typescript is acceptable for the printer; in addition, submit a fully legible carbon, mimeographed, or Xerox copy. Double space entire manuscript, including references, legends, footnotes, and tables.

PREPARATION OF ILLUSTRATIONS Label all illustrations (photographs, micrographs, charts, or drawings) sequentially with arabic numerals; designate them as figures. Number tables with roman numerals. In the margin of the manuscript, indicate approximately where each figure or table should appear.

PAGE SIZE The text block on each Journal page is 5 × 7<sup>3</sup>/<sub>4</sub> inches. Plan illustrations, including their legends, to fit within these dimensions after reduction.

LINE DRAWINGS Provide original ink drawing (preferably  $8\frac{1}{2} \times 11$  inches maximum). These give the sharpest reproductions. The lettering should be sans serif, and of consistent size, and so planned that the size of the lettering is in scale with the size and complexity of the figures. Lettering and labeling should be large enough to permit reduction to a minimum size. Capital letters and numbers should be  $1\frac{1}{2}$  to  $1\frac{3}{4}$  mm high after reduction.

MICROGRAPHS Micrographs should be planned so that they can be reproduced same size (not reduced). Figure numbers and lettering should be  $2\frac{1}{2}$  to 3 mm high in sans serif style. If authors are unable to provide figures labeled satisfactorily, the Press will label them; in this case, the author must provide an overlay or diagram to show exactly where such labeling should appear.

In general, limit the field of a micrograph to the regions and structures discussed in the report; eliminate irrelevant structures or expanses. Cropping unnecessary areas from micrographs saves space and expense. If a micrograph must be specifically oriented, mark "top" on the back of the print.

Micrographs that are grouped should be of uniform size and shape, with the edges of the prints touching. The engraver will tool thin white lines between figures. Plan the over-all size of the grouping so the legend will fit below the composite; if the grouping occupies a full page, the legend will appear at the foot of the facing page. The Journals Office prefers that multiple photographs be submitted unmounted with a diagram showing the desired layout.