## Volume 163 No. 4 April 1, 1986

- 759 R. EPSTEIN, G. SHAM, J. WOMACK, J. YAGÜE, E. PALMER, and M. COHN. The cytotoxic T cell response to the male-specific histocompatibility antigen (H-Y) is controlled by two dominant immune response genes, one in the MHC, the other in the Tarα-locus
- 774 RICHARD P. ARTHUR and DON MASON. T cells that help B cell responses to soluble antigen are distinguishable from those producing interleukin 2 on mitogenic or allogeneic stimulation
- 787 CHANG-YUIL KANG and HEINZ KOHLER. Immunoglobulin with complementary paratope and idiotope
- 797 CARL WALTENBAUGH, LIZHE SUN, and HUAN-YAO LEI. Regulation of immune responses by I-J gene products. VI. Recognition of I-E molecules by I-J-bearing suppressor factors
- REINHOLD E. SCHMIDT, GAIL T. BARTLEY, SHINJI S. LEE, JOHN F. DALEY, HANS DIETER ROYER, HERBERT LEVINE, ELLIS L. REINHERZ, STUART F. SCHLOSSMAN, and JEROME RITZ. Expression of the NKTa clonotype in a series of human natural killer clones with identical cytotoxic specificity
- 826 R. JOHN LOONEY, DANIEL H. RYAN, KAZUHIKO TAKAHASHI, HOWARD B. FLEIT, HARVEY J. COHEN, GEORGE N. ABRAHAM, and CLARK L. ANDERSON. Identification of a second class of IgG Fc receptors on human neutrophils. A 40 kilodalton molecule also found on eosinophils
- 837 TSUKASA SEYA, JERRY R. TURNER, and JOHN P. ATKINSON. Purification and characterization of a membrane protein (gp45-70) that is a cofactor for cleavage of C3b and C4b
- 856 JUN-ICHI MIYAZAKI, ETTORE APPELLA, HONG ZHAO, JAMES FORMAN, and KEIKO OZATO. Expression and function of a nonglycosylated major histocompatibility class I antigen
- 872 WILLIAM E. BOWERS and MARY R. BERKOWITZ. Differentiation of dendritic cells in cultures of rat bone marrow cells
- 884 S. M. Wahl, D. A. Hunt, J. B. Allen, R. L. Wilder, L. Paglia, and A. R. Hand. Bacterial cell wall-induced hepatic granulomas. An in vivo model of T cell-dependent fibrosis
- 903 Lynda A. Morrison, Aron E. Lukacher, Vivian L. Braciale, David P. Fan, and Thomas J. Braciale. Differences in antigen presentation to MHC class I- and class II-restricted influenza virus-specific cytolytic T lymphocyte clones
- 922 A. Granelli-Piperno, L. Andrus, and R. M. Steinman. Lymphokine and nonlymphokine mRNA levels in stimulated human T cells. Kinetics, mitogen requirements, and effects of cyclosporin A
- 938 JAMES R. DASCH and PATRICIA P. JONES. Independent regulation of IgM, IgD, and Ia antigen expression in cultured immature B lymphocytes
- 952 PENTTI UKKONEN, VICTORIA LEWIS, MARK MARSH, ARI HELENIUS, and IRA MELL-MAN. Transport of macrophage Fc receptors and Fc receptor-bound ligands to lysosomes
- 972 NOBUKATA SHINOHARA, JEFFREY A. BLUESTONE, and DAVID H. SACHS. Cloned cytotoxic T lymphocytes that recognize an I-A region product in the context of a class I antigen
- 981 GEORG KRAAL, MATTHIS BREEL, MARGA JANSE, and GERRY BRUIN. Langerhans' cells, veiled cells, and interdigitating cells in the mouse recognized by a monoclonal antibody
- 998 JONATHON SPRENT, MARY SCHAEFER, DAVID Lo, and ROBERT KORNGOLD. Properties of purified T cell subsets. II. In vivo responses to class I vs. class II H-2 differences

## BRIEF DEFINITIVE REPORTS

- 1012 Prakash D. Shah, Jan Keij, Susan M. Gilbertson, and Donald A. Rowley. Thy-1<sup>+</sup> and Thy-1<sup>-</sup> natural killer cells suppress dendritic cells
- 1018 DAVID B. LEWIS, ALF LARSON, and CHRISTOPHER B. WILSON. Reduced interferon-gamma mRNA levels in human neonates. Evidence for an intrinsic T cell deficiency independent of other genes involved in T cell activation
- 1024 Frank Emmrich, Jelle Thole, Jan van Embden, and Stefan H. E. Kaufmann. A recombinant 64 kD protein of *Mycobacterium bovis* Bacillus Calmette-Guerin specifically stimulates human T4 clones reactive to mycobacterial antigens

1030 KAYO INABA, MAKOTO KITAURA, TAKUMA KATO, YOSHIHIKO WATANABE, YOSHIMA KAWADE, and SHIGERU MURAMATSU. Contrasting effect of  $\alpha/\beta$ - and  $\gamma$ -interferons on expression of macrophage Ia antigens

## COPYEDITING AND PRODUCTION Samuel K. Faith and Vera J. Appleyard

PREPARATION OF MANUSCRIPT Articles should conform to the style of a current issue of this journal or to the recommendations of the Council of Biology Editors Style Manual (5th edition, 1983, Council of Biology Editors, Inc., 9650 Rockville Pike, Bethesda, MD 20814). Note that each reference should contain the title of the pertinent citation. Please supply a brief title for a running head, not exceeding 55 characters and spaces, with no abbreviations. Capitalize trade names and give manufacturers' full names and addresses. Only original typescript is acceptable for the printer; in addition, submit a fully legible carbon, mimeographed, or Xerox copy. Double space entire manuscript, including references, legends, footnotes, and tables.

PREPARATION OF ILLUSTRATIONS Label all illustrations (photographs, micrographs, charts, or drawings) sequentially with arabic numerals; designate them as figures. Number tables with roman numerals. In the margin of the manuscript, indicate approximately where each figure or table should appear.

PAGE SIZE The text block on each Journal page is  $5 \times 7^{3}$ 4 inches. Plan illustrations, including their legends, to fit within these dimensions after reduction.

LINE DRAWINGS Provide original ink drawing (preferably  $8\frac{1}{2} \times 11$  inches maximum). These give the sharpest reproductions. The lettering should be sans serif, and of consistent size, and so planned that the size of the lettering is in scale with the size and complexity of the figures. Lettering and labeling should be large enough to permit reduction to a minimum size. Capital letters and numbers should be  $1\frac{1}{2}$  to  $1\frac{3}{4}$  mm high after reduction.

MICROGRAPHS Micrographs should be planned so that they can be reproduced same size (not reduced). Figure numbers and lettering should be  $2\frac{1}{2}$  to 3 mm high in sans serif style. If authors are unable to provide figures labeled satisfactorily, the Press will label them; in this case, the author must provide an overlay or diagram to show exactly where such labeling should appear.

In general, limit the field of a micrograph to the regions and structures discussed in the report; eliminate irrelevant structures or expanses. Cropping unnecessary areas from micrographs saves space and expense. If a micrograph must be specifically oriented, mark "top" on the back of the print.

Micrographs that are grouped should be of uniform size and shape, with the edges of the prints touching. The engraver will tool thin white lines between figures. Plan the over-all size of the grouping so the legend will fit below the composite; if the grouping occupies a full page, the legend will appear at the foot of the facing page. The Journals Office prefers that multiple photographs be submitted unmounted with a diagram showing the desired layout.