

THE JOURNAL OF EXPERIMENTAL MEDICINE

Volume 195, No. 2, January 21, 2002

Contents

Commentary

- F5** *Michael E. Lamm and Julia M. Phillips-Quagliata.* Origin and homing of intestinal IgA anti-body-secreting cells

Articles

- 151** *Satoshi Takaki, Hatsue Morita, Yoshinari Tezuka, and Kiyoshi Takatsu.* Enhanced hematopoiesis by hematopoietic progenitor cells lacking intracellular adaptor protein, Lnk
- 161** *Kazuyoshi Takeda, Mark J. Smyth, Erika Cretney, Yoshihiro Hayakawa, Nobuhiko Kayagaki, Hideo Yagita, and Ko Okumura.* Critical role for tumor necrosis factor-related apoptosis-inducing ligand in immune surveillance against tumor development
- 171** *Jeffrey P. Walterscheid, Stephen E. Ullrich, and Dat X. Nghiem.* Platelet-activating factor, a molecular sensor for cellular damage, activates systemic immune suppression
- 181** *Yijin Li, Hui Li, and Martin Weigert.* Autoreactive B cells in the marginal zone that express dual receptors
- 189** *Kazunori Inabe, Masamichi Ishiai, Andrew M. Scharenberg, Norman Freshney, Julian Downward, and Tomohiro Kurosaki.* Vav3 modulates B cell receptor responses by regulating phosphoinositide 3-kinase activation
- 201** *Nacksung Kim, Masamichi Takami, Jaerang Rho, Regis Josien, and Yongwon Choi.* A novel member of the leukocyte receptor complex regulates osteoclast differentiation
- 211** *V. Michael Holers, Guillermina Girardi, Lian Mo, Joel M. Guthridge, Hector Molina, Silvia S. Pierangeli, Ricardo Espinola, Liu E. Xiaowei, Dailing Mao, Christopher G. Vialpando, and Jane E. Salmon.* Complement C3 activation is required for antiphospholipid antibody-induced fetal loss
- 221** *Peter McGuirk, Chantelle McCann, and Kingston H.G. Mills.* Pathogen-specific T regulatory 1 cells induced in the respiratory tract by a bacterial molecule that stimulates interleukin 10 production by dendritic cells: a novel strategy for evasion of protective T helper type 1 responses by *Bordetella pertussis*
- 233** *Elizabeth Alcamo, Nir Hacohen, Leah C. Schulte, Paul D. Rennert, Richard O. Hynes, and David Baltimore.* Requirement for the NF- κ B family member RelA in the development of secondary lymphoid organs
- 245** *Norbert Gerdes, Galina K. Sukhova, Peter Libby, Rebecca S. Reynolds, James L. Young, and Uwe Schönbeck.* Expression of interleukin (IL)-18 and functional IL-18 receptor on human vascular endothelial cells, smooth muscle cells, and macrophages: implications for atherogenesis

Contents continued

Cover picture: The role of the RelA in the development of secondary lymphoid organs has not been possible to determine as RelA-deficient mice die at embryonic day 15. However, mice deficient for both RelA and TNFR1 are viable and reveal an essential role for RelA in the development of lymph nodes, Peyer's patches, and spleen. The cover images show the microarchitecture of spleens from wild-type (left) and TNFR1/RelA-deficient (right) mice. TNFR1/RelA mutant spleens have severely reduced T and B cell zones (top: CD3 in purple and B220 in brown to mark T and B cell zones, respectively), and lack follicular dendritic cells (bottom: CR1 in purple) and germinal centers (bottom: PNA in brown). See related article by Alcamo et al., pp. 233–244.

- 259 ***Ying Liu, Pedram Hamrah, Qiang Zhang, Andrew W. Taylor, and M. Reza Dana.*** Draining lymph nodes of corneal transplant hosts exhibit evidence for donor major histocompatibility complex (MHC) class II-positive dendritic cells derived from MHC class II-negative grafts
- 269 ***Edward P. Bowman, Nelly A. Kuklin, Kenneth R. Youngman, Nicole H. Lazarus, Eric J. Kunkel, Junliang Pan, Harry B. Greenberg, and Eugene C. Butcher.*** The intestinal chemokine thymus-expressed chemokine (CCL25) attracts IgA antibody-secreting cells

Brief Definitive Report

- 277 ***Roseann Waterhouse, Cam Ha, and Gabriela S. Dveksler.*** Murine CD9 is the receptor for pregnancy-specific glycoprotein 17