

THE JOURNAL OF EXPERIMENTAL MEDICINE

Volume 197, No. 12, June 16, 2003

Contents**Articles**

- 1603** *Nicole J. Horwood, Tara Mahon, John P. McDaid, Jamie Campbell, Hiroyuki Mano, Fionula M. Brennan, David Webster, and Brian M.J. Foxwell.* Bruton's tyrosine kinase is required for lipopolysaccharide-induced tumor necrosis factor α production
- 1613** *Vallabhapurapu Sivakumar, Kirsten J.L. Hammond, Norma Howells, Klaus Pfeffer, and Falk Weil.* Differential requirement for Rel/nuclear factor κ B family members in natural killer T cell development
- 1623** *Dirk Elewaut, Raziya B. Shaikh, Kirsten J.L. Hammond, Hilde De Winter, Andrew J. Leishman, Stephane Sidobre, Olga Turovskaya, Theodore I. Prigozy, Lisa Ma, Theresa A. Banks, David Lo, Carl F. Ware, Hilde Cheroutre, and Mitchell Kronenberg.* NIK-dependent RelB activation defines a unique signaling pathway for the development of V α 14i NKT cells
- 1635** *Elmar Jaeckel, Ludger Klein, Natalia Martin-Orozco, and Harald von Boehmer.* Normal incidence of diabetes in NOD mice tolerant to glutamic acid decarboxylase
- 1645** *Naglaa H. Shoukry, Arash Grakoui, Michael Houghton, David Y. Chien, John Ghayeb, Keith A. Reimann, and Christopher M. Walker.* Memory CD8 $^{+}$ T cells are required for protection from persistent hepatitis C virus infection
- 1657** *Joseph B. El Khoury, Kathryn J. Moore, Terry K. Means, Josephine Leung, Kinya Terada, Michelle Toft, Mason W. Freeman, and Andrew D. Luster.* CD36 mediates the innate host response to β -amyloid
- 1667** *Madhav V. Dhodapkar, Matthew D. Geller, David H. Chang, Kanako Shimizu, Shin-Ichiro Fujii, Karita M. Dhodapkar, and Joseph Krasovsky.* A reversible defect in natural killer T cell function characterizes the progression of premalignant to malignant multiple myeloma
- 1677** *Kathryn A. Pape, Valerie Kouskoff, David Nemazee, H. Lucy Tang, Jason G. Cyster, Lina E. Tze, Keli L. Hippen, Timothy W. Behrens, and Marc K. Jenkins.* Visualization of the genesis and fate of isotype-switched B cells during a primary immune response
- 1689** *Chang-Hung Chen, Carole Seguin-Devaux, Nancy A. Burke, Timothy B. Oriss, Simon C. Watkins, Neil Clipstone, and Anuradha Ray.* Transforming growth factor β blocks Tec kinase phosphorylation, Ca $^{2+}$ influx, and NFATc translocation causing inhibition of T cell differentiation
- 1701** *Petronela Ancuta, Ravi Rao, Ashlee Moses, Andrew Mehle, Sunil K. Shaw, F. William Luscinskas, and Dana Gabuzda.* Fractalkine preferentially mediates arrest and migration of CD16 $^{+}$ monocytes
- 1709** *Remy Bosselut, Terry I. Guinter, Susan O. Sharow, and Alfred Singer.* Unraveling a revealing paradox: why major histocompatibility complex I–signaled thymocytes “paradoxically” appear as CD4 $^{+}$ 8 $^{\text{lo}}$ transitional cells during positive selection of CD8 $^{+}$ T cells

Contents continued

Cover picture: Visualization of isotype switching in the spleen, 4 d after immunization. The image shows antigen-specific, isotype-switched B cells (green) coexisting in an early germinal center (the central area containing red B220 $^{+}$, IgD $^{-}$ cells surrounded by purple B220 $^{+}$, IgD $^{+}$ naive B cells), the marginal zone (the rim area containing red B220 $^{+}$, IgD $^{-}$ cells), and red pulp (the dark area below the B220 $^{+}$ follicle). See related article by Pape et al., pp. 1677–1687.

- 1721 **Xingluo Liu, Jian Xin Gao, Jing Wen, Lijie Yin, Ou Li, Tao Zuo, Thomas F. Gajewski, Yang-Xin Fu, Pan Zheng, and Yang Liu.** B7DC/PDL2 promotes tumor immunity by a PD-1-independent mechanism
- 1731 **Xu Hong, Lu Lei, and Rickard Glas.** Tumors acquire inhibitor of apoptosis protein (IAP)-mediated apoptosis resistance through altered specificity of cytosolic proteolysis
- 1745 **Mingfang Lu, Mei Zhang, Richard L. Kitchens, Susan Fosmire, Akira Takashima, and Robert S. Munford.** Stimulus-dependent deacylation of bacterial lipopolysaccharide by dendritic cells
- 1755 **Peter Vajkoczy, Sabine Blum, Mathias Lamparter, Reinhard Mailhammer, Ralph Erber, Britta Engelhardt, Dietmar Vestweber, and Antonis K. Hatzopoulos.** Multistep nature of microvascular recruitment of ex vivo-expanded embryonic endothelial progenitor cells during tumor angiogenesis
- 1767 **Bernardo Reina-San-Martin, Simone Difilippantonio, Leif Hanitsch, Revati F. Masilamani, André Nussenzweig, and Michel C. Nussenzweig.** H2AX is required for recombination between immunoglobulin switch regions but not for intra-switch region recombination or somatic hypermutation

Brief Definitive Reports

- 1779 **Yuichi Harada, Masamichi Muramatsu, Toshikatsu Shibata, Tasuku Honjo, and Kazumichi Kuroda.** Unmutated immunoglobulin M can protect mice from death by influenza virus infection
- 1787 **Clett Erridge, John Stewart, and Ian R. Poxton.** Monocytes heterozygous for the Asp299Gly and Thr399Ile mutations in the Toll-like receptor 4 gene show no deficit in lipopolysaccharide signalling