- Louis S. Premkumar and Anthony Auerbach. Stoichiometry of recombinant *N*-methyl-D-aspartate receptor channels inferred from single-channel current patterns
- 503 Klaus Schumann, Christoph Romanin, Werner Baumgartner, and Klaus Groschner. Intracellular Ca²⁺ inhibits smooth muscle L-type Ca²⁺ channels by activation of protein phosphatase type 2B and by direct interaction with the channel
- 515 DAVID H. HACKOS and JUAN I. KORENBROT. Calcium modulation of ligand affinity in the cyclic GMP-gated ion channels of cone photoreceptors
- 529 EDWARD J. KAFTAN, BARBARA E. EHRLICH, and JAMES WATRAS. Inositol 1,4,5-Trisphosphate (InsP₃) and calcium interact to increase the dynamic range of InsP₃ receptor-dependent calcium signaling
- 539 JOHN G. STARKUS, LIOBA KUSCHEL, MARTIN D. RAYNER, and STEFAN H. HEINEMANN. Ion conduction through C-type inactivated *Shaker* channels
- 551 Christoph Fahkle, Christine Dürr, and Alfred L. George, Jr. Mechanism of ion permeation in skeletal muscle chloride channels
- 565 M. Yoshino, S.Y. Wang, and C.Y. Kao. Sodium and calcium inward currents in freshly dissociated smooth myocytes of rat uterus
- 579 RICCARDO OLCESE, RAMÓN LATORRE, LIGIA TORO, FRANCISCO BEZANILLA, and ENRICO STE-FANI. Correlation between charge movement and ionic current during slow inactivation in *Shaker* K⁺ channels
- 591 Anthony A. Fodor, Kevin D. Black, and William N. Zagotta. Tetracacine reports a conformational change in the pore of cyclic nucleotide–gated channels
- MARK R. BOWLBY, DEBRA A. FADOOL, TODD C. HOLMES, and IRWIN B., LEVITAN. Modulation of the Kv1.3 potassium channel by receptor tyrosine kinases
- 611 K.S. Kits, J.C. Lodder, and M.J. Veerman. Phe-Met-Arg-Phe-amide activates a novel voltagedependent K⁺ current through a lipoxygenase pathway in molluscan neurones
- 629 EMILIA M. HOGAN, BRUCE A. DAVIS, and WALTER F. BORON. Intracellular Cl⁻ dependence of Na-H exchange in barnacle muscle fibers under normotonic and hypertonic conditions
- 641 Correction